



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/37	A1	(11) International Publication Number: WO 98/39471 (43) International Publication Date: 11 September 1998 (11.09.98)
(21) International Application Number: PCT/US98/03929 (22) International Filing Date: 6 March 1998 (06.03.98) (30) Priority Data: 60/038,940 7 March 1997 (07.03.97) US (71) Applicant: TROPIX, INC. [US/US]; 47 Wiggins Avenue, Bedford, MA 01730 (US). (72) Inventors: BRONSTEIN, Irena; 11 Ivanhoe Street, Newton, MA 02158 (US). VOYTA, John; 99 Maynard Farm Road, Sudbury, MA 01776 (US). PALMER, Michelle; 87 Medford Street, Arlington, MA 02174 (US). TILLOTSON, Bonnie; Unit 1, 105 Beech Street, Belmont, MA 02178 (US). (74) Agents: KELBER, Steven, B. et al.; Oblon, Spivak, McClelland, Maier & Neustadt, P.C., Crystal Square Five, 4th floor, 1755 Jefferson Davis Highway, Arlington, VA 22202 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PROTEASE INHIBITOR ASSAY		
(57) Abstract <p>Heterogenous and homogenous assays are provided for the detection of protease inhibitory activity in a sample or target compound, taking advantage of the chemiluminescent characteristics of 1,2-dioxetanes. In the heterogenous assay, a peptide bearing a cleavage site for the protease of interest is provided with a first member of a first ligand binding pair at one end, and a first member of a second ligand binding pair at the other end. The other member of the first ligand binding pair is attached to a surface, which binds the peptide, or protease substrate, to the surface. The peptide substrate is combined with the protease and target compound or sample. Substrate cleavage, if not inhibited, is allowed to occur, and any unbound cleaved fragments are removed. An enzyme complexed with the second member of the second ligand binding pair is added, and allowed to bind to any of the (uncleaved) first member of the second ligand binding pair remaining. Unbound complex is removed, and a 1,2-dioxetane substrate for the enzyme is added. If any peptide substrate has not been cleaved, the dioxetane will chemiluminesce, indicating inhibitory activity. In a homogenous assay, the same substrate bears at one end a fluorescent energy accepting moiety, and at the other end a 1,2-dioxetane or precursor. If the substrate is cleaved by the protease, the dioxetane and the fluorescent moiety are not in close physical relationship, and no energy transfer occurs when the dioxetane is caused to decompose. If cleavage has not occurred, indicating inhibition, when the dioxetane is caused to decompose, energy is transferred to the fluorescing entity, which releases light of a wavelength recognizably distinct from that of the dioxetane.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TITLE OF THE INVENTION**PROTEASE INHIBITOR ASSAY**

This application is a regular National application claiming priority from Provisional Application. U.S. Application Serial No. 60/038,940 filed March 7, 1997.

BACKGROUND OF THE INVENTION**Field of the Invention**

This invention pertains to the use of chemiluminescent 1,2-dioxetanes in homogenous or heterogeneous assays to detect proteases inhibitors.

Discussion of the Background

The identification of novel therapeutics that block or inhibit inimical proteases, or proteases that mediate disease conditions, such as the 11-kd protease encoded by the human immunodeficiency virus 1 (HIV-1) is a key step in slowing the disease process of AIDS. Retroviral proteases are essential in the process of viral gag-pol polyproteins of the HIV-1 and HIV-2 viruses. There are a few highly conserved consensus sequences in retroviral polyproteins, one of which consists of a pentapeptide (Ser/Thr)-X-X'-(Tyr/Phe)-Pro. Cleavage occurs between the Tyr or Phe and Pro residues. Blocking activity of these proteases will interfere with the progression of HIV infection. Although potent drugs which block HIV protease activity have been found, there is an ongoing need to find and develop novel inhibitors.

Current methods utilized in rapid screening of protease inhibitors are subject to many interferences from a variety of sources. The most common non-isotopic approach is a fluorescent assay. In one case, such as in the detection of HIV protease, the fluorescent

substrate is labeled with a fluorescent dansyl group on one end of a peptide and a quencher on the other. An increase in fluorescence signal occurs upon cleavage of the protease due to the fact that the emitter and the quencher are separated as described in Matayoshi et al., 1990, Science 247: 954-958. Fluorescent substrates for other proteases can be designed with a terminal fluorophore which emits a fluorescent signal upon cleavage by the enzyme.

Both of the above assay approaches are commonly used as high throughput assays for screening large chemical, natural product and combinatorial libraries. These assays tend to have problems related to autofluorescence of biological components due to the nature of the molecules which are screened. Many of the compounds and natural product extracts are colored or fluorescent and are present in the solution when the assay signal is monitored. This results in an assay interference which limits the detection sensitivity and the dynamic range of the assay. This interference can easily be interpreted as an inhibition of the enzyme, making it difficult to determine true positive inhibition, thus, requiring extensive follow-up assays to distinguish true positives from the false positives.

U.S. Patent 5,591,591, assigned to Tropix, describes assays for the detection of proteases wherein a dioxetane compound bearing a proteolytic enzyme-specific amino acid or peptide, is added to a sample suspected of containing the protease, and the amino acid is removed by enzymatic reaction, causing the dioxetane to decompose and chemiluminescence.

SUMMARY OF THE INVENTION

An alternative approach to the above homogeneous approaches is to use homogenous or heterogeneous assays which are not subject to interference. The present inventors have developed highly sensitive assays using a chemiluminescent 1,2-dioxetane

substrate for high throughput screening of HIV-1 protease activity. The present invention provides the advantage of an assay that it is not subject to interferences from colored or fluorescent compounds, and therefore is more sensitive and exhibits a greater dynamic range compared to a direct, fluorescent enzyme assay. This assay, adapted for the detection of HIV protease inhibitors, utilizes a synthetic peptide substrate I (Fam-spacer-Ser-Gln-Asn-Try-Pro-Ile-Val-Gln-spacer-(Biotin)-NH₂) whose sequence is derived from the native cleavage site of the Gag polyprotein. (Fam is used herein to indicate fluorescein). Many high throughput HIV screening methodologies exist which utilize large quantities of reagents and involve more laborious manipulations. The assay of the present invention may be advantageously formatted as a simple single plate endpoint assay which is sensitive down to fmoles of captured peptide which is particularly useful for high throughout screening, although it may also be presented as a conventional two-step transfer and dilution capture assay. The assay uses a chemiluminescent 1,2-dioxetane alkaline phosphatase substrate in an immunoassay format for the sensitive detection of cleaved peptide. The uncleaved peptide is recognized by an anti-fluorescein alkaline phosphatase conjugated antibody. Capture conditions have been optimized to assure a linear response to cleaved peptide concentration. This response correlates well to HPLC analysis of cleaved product. The HIV assay has been validated for acetyl-pepstatin, a known HIV-1 inhibitor. This robust assay is amenable to automation and can be used to screen large numbers of compounds in a cost effective 96 well format.

In an alternative embodiment, a homogenous assay less sensitive to color or fluorescence interference than prior art assays is used. In this assay, the same peptide is employed, but labeled at one terminus with fluorescein or other fluorescing energy

acceptor, and at the other end by a 1,2-dioxetane label covalently attached to the peptide terminus. After admixture of the target compound, the dioxetane is caused to decompose by addition of a chemical (enzymatic or non-enzymatic) or another trigger, such as application of heat or change in pH. If the peptide has not been cleaved, the dioxetane is in close physical relationship with the fluorescent energy acceptor, and upon triggering, energy transfer assisted fluorescence is observed. The light has a characteristic color dependent on the fluorescent emitter, such as green for fluorescein. If the peptide has been cleaved, the chemiluminescent light of the dioxetane itself, typically a bluer light, is observed.

DETAILED DESCRIPTION OF THE DRAWINGS

Figure 1. Illustration of two labeled HIV-1 substrates I and II whose sequences are derived from a natural processing site for aspartyl HIV-1 protease. Incubation of these two HIV-1 peptides with rec HIV-1 protease results in a specific cleavage between the Try-Pro bond as reported in Kohl, NE et al.(1988), Proc. Acad. Sci. USA, 85, 4684 and Billich, SW, et al. (1988), J. Biol. Chem. 263, 17905. These are compared against the prior art substrate as reported in Mayatoshi et al.

Figure 2. HIV-1 Protease Assay. Steps include 1. Addition of rec HIV-1 protease, acetyl-pepstatin(3) and/or screening compound to NeutrAvidin coated plates, with a preincubation. 2. Addition of HIV-1 protease peptide I followed with a 60 min. Incubation at 37 °C. 3. Wash out cleaved peptide fragment. 4. Add anti-fluorescein alkaline phosphatase antibody to detect uncleaved bound peptide and then 5. Add CSPD/Sapphire II and measure emitted light in a TR717 luminometer(Tropix).

Figure 3. The phosphate group on the CSPD substrate is cleaved by alkaline phosphatase to generate the unstable anion intermediate 1. Intermediate 1 then undergoes decomposition producing a long-lived emission at 470nm.

Figure 4. HIV-1 protease peptide I titration on Neutravidin coated plates. 100ul of diluted peptide was added to various densities of coated Neutravidin wells. Plates were washed and peptide was detected with an anti-fluorescein alkaline phosphatase conjugate followed with the addition of CSPD/Sapphire II. Light emission was measured in a TR717 microplate luminometer(Tropix).

Figure 5(A&B). Comparison of HIV-1 peptide I cleavage by rec HIV-1 protease by HPLC analysis and capture on Neutravidin coated plates. 35nM of rec HIV-1 protease was added to 10uM HIV peptide in eppendorf tubes and incubated at 37 °C. At specific time points 0.1% TFA was added to stop the reaction and tubes placed on ice. Reaction products were analyzed by HPLC. Injection volumes were 100ul and absorbency was monitored @ 490 nm. The percentage of cleaved product was calculated from the peak areas. For peptide capture experiments, 100ul of a 1:50 dilution was added to Neutravidin (5ug/ml) coated plates and incubated at ambient temp. for 60 min. after a wash step, captured peptide was detected with anti-fluorescein conjugated alkaline phosphatase antibody followed by another wash step and then the addition of CSPD/Sapphire II. Light emission was measured in a TR717 microplate luminometer(Tropix).

Figure 6. Hydrolysis of EDANS/DABCYL HIV-1 protease Prior Art Substrate by rec HIV-1 protease. Fluorogenic HIV-1 peptide II was added to protease buffer pH 4.7 in 96 well plates. Rec HIV-1 protease was added at ambient temperature and immediately read

over a 105 minute time period on a LS50B fluorescence spectrophotometer (Perkin-Elmer). Excitation was set at 340nm and emission was set at 490nm.

Figure 7. Hydrolysis and capture of HIV-1 protease substrate I on Neutravidin coated plates. Dilutions of rec HIV-1 protease diluted in protease buffer pH 5.5 were added to wells previously coated with 0.1ug/ml Neutravidin. 10 pmoles of protease substrate was added to start the reaction and the plate was incubated for 1 hour at 37 °C. After a wash step, captured peptide was detected with anti-fluorescein conjugated alkaline phosphatase antibody followed by another wash step and then the addition of CSPD/Sapphire II. Light emission was measured in a TR717 microplate luminometer(Tropix). 100% cleavage of the substrate is not obtained for substrate 1. The date has been adjusted to reflect this phenomenon. In actuality, the detected background may be different.

Figure 8. Rec HIV-1 protease inhibition by acetyl-pepstatin in a HIV-1 capture assay. 2.5 pmoles of rec HIV-1 protease diluted in protease buffer pH 5.5 was added to Neutravidin (0.1ug/ml) coated wells. Dilutions of acetyl-pepstatin in DMSO were added to wells followed with a preincubation of 5 minutes. 20 pmoles of HIV-1 protease substrate was added and the plate incubated for 1 hour at 37 °C. After a wash step captured peptide was detected with an anti-fluorescein conjugated alkaline phosphatase antibody. After a second wash step, CSPD/Sapphire II was added and light emission was measured in a TR717 microplate luminometer(Tropix).

Figure 9. Screening compound plate-1001 at 25uM (5% DMSO) for rec HIV-1 protease inhibition. 2.5 pmoles of rec HIV-1 protease diluted in protease buffer pH 5.5 was added to Neutravidin coated plates. 5ul from a compound screening plate largely composed of drug standards was added and preincubated for 5 minutes. 20 pmoles of HIV-

1 protease substrate I was added and the plate was incubated for 60 minutes at 37 °C. After a wash step, captured peptide was detected with anti-fluorescein conjugated alkaline phosphatase antibody. After a second wash step, CSPD/Sapphire II was added and light emission was measured in a TR717 microplate luminometer(Tropix).

Figure 10. HIV FLAG Peptide titration on Neutravidin and Anti-FLAG MI

monoclonal antibody coated plates. 100ul of diluted HIV FLAG peptide was added to various densities of pre-coated Neutravidin or Anti FLAG MI monoclonal antibody coated plates. Coating was achieved by diluting a given amount of Neutravidin or Anti-FLAG MI monoclonal antibody and BSA in PBS so as to achieve a total protein concentration of 5 µg/ml. Plates were incubated for 1 hour at 37 °C followed by a 3x wash with TBS/0.05% Tween/3mM CaCl₂. 100 ul of a 1:20,000 dilution of Avidix-AP was added and incubated for 1 hour at room temp. After 3 washes in TBS/Tween/CaCl₂ buffer and 1 wash with 1x Tris/1mM MgCl/3mM CaCl pH 9.8, 100 ul ready to use CSPD/Sapphire II was added and wells incubated for 30 min at room temp. Luminescence was then measured in a TR717 microplate luminometer (Tropix). The present inventors have discovered novel peptide capture conditions which permit the assay to be performed as a single well (plate) assay as opposed to a 2 step transfer and dilution assay. As shown in Figure 12, the signals are relatively the same when done as a one step or two step assay suggesting that non-specific cleavage of other assay components by the rec HIV-1 protease is nonexistent.

Figure 11. Capture of HIV FLAG and HIV FAM Peptides on Neutravidin (5 ug/ml) coated plates post hydrolysis with rec HIV-1 protease (50nM). 50 nM rec HIV-1 protease was added to 1uM HIV FLAG or HIV FAM peptides in protease buffer (0.1mM NaAcetate, 1M NaCl, 1mM EDTA, 1mM DTT, 1 mg/ml BSA, pH 4.7). Biotin at 0.5 mM

and 5 Units of enterokinase were added to control tubes. Reactions were done for 1 hour at 37 °C in eppendorf tubes. All tubes were then placed on ice to stop the reaction. An additional 50nM rec HIV-1 protease was added to 2 of the reaction tubes and incubated further for 30 minutes at 37 °C. HIV FLAG peptide containing tubes were diluted 1:100 (10nM final) and HIV FAM peptide tubes were diluted 1:100,000 (0.01nM final) in protease buffer. 100ul was added to Neutravidin coated wells (5 ug/ml) and incubated for 1 hour at room temp. Wells were then washed with 3x PBS/Tween/CaCl buffer. 100ul of 1:5000 dilution of Anti FITC-AP Fab Fragment was added to HIV FAM peptide containing wells and 100ul of a precomplexed Anti-FLAG MI antibody (1:1000) and Goat anti mouse-AP conjugate (1:10,000) was added to HIV FLAG peptide containing wells. Plates were incubated for additional hour at room temp then washed 3x in TBS/Tween/CaCl and 1x Tris/MgCl/CaCl pH 9.8. 100ul ready to use CSPD/Sapphire II was added to wells and incubated 30 minutes at room temp. Luminescence was then measured in a TR717 microplate luminometer.

Figure 12. Titration of rec HIV-1 Protease I with 100nM HIV FLAG Peptide. Diluted rec HIV-1 protease was added to 100nM HIV FLAG peptide in protease buffer. Reactions were done either in eppendorf tubes or directly in wells precoated with Anti-FLAG MI monoclonal antibody (0.1 ug/ml). Competing FLAG octapeptide at 2.5uM was added to control tubes and wells. Incubations were done for 1 hour at 37 °C, then eppendorf reaction products transferred to wells, and further incubated for 1 hour at room temp. Plates were washed 3x in PBS/Tween/CaCl buffer followed with addition of a 1:20,000 dilution of Avidix-AP. Plates were further incubated for 1 hour at room temp and then washed 3x with PBS/Tween/CaCl buffer and 1x with Tris/MgCl/CaCl pH 9.8 buffer.

100ul CSPD/Sapphire II was added and wells incubated for 30 minutes to room temp.

Luminescence was then measured in a TR717 microplate luminometer (Tropix).

Figure 13. Inhibition of 12.5 nM rec HIV-1 Protease with Acetyl-Pepstatin. Assays consisted of reactions with 100 nM HIV FLAG Peptide in wells coated with Anti-FLAG MI antibody (0.1 ug/ml) for 1 hour at 37 °C. Dilutions of acetyl-pepstatin were preincubated for 5 minutes with rec HIV-1 enzyme before adding to HIV FLAG peptide substrate in protease buffer. All additions were made on a Zymark Rapid Plate. The plate was then washed 4x with TBS/3mM CaCl₂/0.05% Tween on Tecan 96 plate washer. 100ul of a 1:20,000 dilution of Avidix-AP conjugate was added to each well and incubated for 1 hour @ room temp. The plate was then washed 3x with above buffer and 1x with Tris/MgCl/CaCl pH 9.8. 100ul CSPD/Sapphire II was added, incubated for 30 min and then light emission was measured in a TR717 luminometer (Tropix). Note: Acetyl-pepstatin is a well known aspartic proteinase inhibitor and is a reported inhibitor of HIV-1 protease (*Pro. Natl. Sci. U.S.A.* 85, 66123, (1988)).

Figure 14. Synthetic peptide for homogenous assay. This peptide, also adapted for HIV-1 protease inhibitor detection, is derivatized at both ends of the molecule, in fashion similar to that employed in the heterogenous assay. In this case, however, one terminus, preferably the carboxy terminus, is labeled with an energy accepting fluorescent moiety, such as fluorescein, and the amino terminus is labeled with a 1,2-dioxetane moiety precursor (14A) which can be photooxygenated in situ to the dioxetane (14B). The dioxetane may be triggered by other than chemical (enzymatic or non-enzymatic) means if necessary.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

A peptide substrate is synthesized which contains the appropriate cleavage site for the target protease. This peptide is labeled, in a heterogenous assay, with one member of a first ligand binding pair, such as biotin, on one end and a member of a second ligand binding pair, such as fluorescein, at the other end. This peptide is then incubated with the protease and a compound of interest to be screened for inhibitory activity, in a well or other solid phase coated with the second binding ligand of said first ligand binding pair, such as avidin or strepavidin.

The spacer between the ligand binding pair members and the peptide of the inventive substrate may be a covalent bond or any covalent binding moiety which does not interfere with either the first or second ligand binding pair or substrate cleavage. Among exemplary sequences are C1-12 alkyls, alkylamino's, carboxylic acids, or any neutral moiety terminating in coupling functionalities at either end. The spacer may be provided with water-solubility enhancing substituents (e.g., carboxy, sulfoxy, halo, etc.) or, where necessary for enzyme/antibody presentation, to cyclize the substrate, bridging agents.

After incubation, the wells are washed, incubated with the second binding member of said second binding ligand pair conjugated with an enzyme which is an effective trigger for a 1,2-dioxetane, such as alkaline phosphatase, washed, incubated with a 1,2-dioxetane substrate such as chlorine substituted phosphate dioxetane (CSPD) and the signal is measured. Higher signals are detected in the presence of an inhibitor.

The 1,2-dioxetanes used as substrates may be any of those described in any of Tropix' prior patents, including 4,931,223; 4,931,569; 4,952,707; 4,956,477; 4,978,614; 5,032,381; 5,112,960; 5,154,772; 5,220,005; 5,225,584; 5,326,882; 5,330,900;

5,336,596; 5,543,295; 5,582,980; 5,605,795; 5,625,007; 5,654,154; and 5,679,803, which are incorporated by reference herein. The above mentioned patents disclose 1,2-dioxetanes as chemiluminescent compounds which can be used as reporter molecules in ultrasensitive assays that can be conducted quickly without resort to exotic conditions or elaborate apparatus, for the detection of a variety of biological molecules. A preferred substrate is CSPD, the chlorine-substituted counterpart of AMPPD, which is the disodium salt of 3-(4-methoxy-spiro[1,2-dioxetane-3,2'-tricyclo[3.3.1.1^{3,7}]decan]4-yl)phenyl phosphate.

The dioxetanes contain a fluorescent chromophore group, preferably an aryl group such as phenyl or naphthyl and an enzyme cleavable group, e.g., a phosphate ester, which when cleaved by the appropriate enzyme, e.g., alkaline phosphatase, forms a negatively charged substituent (e.g., an oxyanion). This destabilizes the dioxetane, thereby causing the dioxetane to decompose to form two carbonyl-containing groups accompanied by the release of light.

To enhance the chemiluminescent signal, and improve signal/noise ratio to permit discrimination between background signals and positive target-responsive signals at very low levels, a water-soluble enhancement agent may be added to the sample prior to or concomitant with the introduction of the dioxetane. Specific enhancement agents which may be used include: quaternary onium polymeric salts such as polyvinylbenzyltributylammonium chloride (Sapphire II, Tropix) or any of those disclosed in U.S. Patent 5,336,596 as potential membrane coatings, neutral detergents such as Tween-20 (Sigma), cationic detergents, such as cetyltrimethylammonium chloride (CTAB, Sigma) and combinations thereof.

Families of proteolytic enzymes which may be targets for determination of inhibitory agents include

Cysteine Proteases

Caspases 1, 2, 3, 6, 7, 8
Cathepsins (B, H, S and L)
Hydrolase
L-proteinases
Calpain
Interleukin converting proteases (ICE)

Serine Proteases

Urokinase
Trypsin
Thrombin
Cathepsin G

Aspartic Proteases

HIV-1 and 2
Yapsin I and YAP 3
Plasmepsin I and II
Cathepsin D and E

Metalloproteinases

Collagenase
Gelatinase A and B
Stromelysin
Aminopeptidase
Elastase

or any of those mentioned in U.S. Patent 5,591,591. Assay conditions and applications for specific proteases are also listed in U.S. Patent 5,591,591. It is possible to use any peptide which is recognized by the protease of interest. Examples of specific peptides and proteases are listed in U.S. Patent 5,591,591. In particular, a peptide is prepared which features a cleavage site for which the target protease is specific. The resulting peptide is short, and easily prepared using conventional synthetic technology.

A large variety of ligand binding pairs can be used for both the first and second ligand binding pair employed. Among preferred labels for either end of the peptide are: Biotin, Fluorescein (FAM), FLAG, HIS tag (6 histidine amino acid sequence), and Digoxin (digoxigenin labeled peptide). These are bound by the other member of the first pair, preferably bound or attached to a solid phase so as to remain through washing and addition of the other member of the second pair, added after an opportunity for cleavage to occur, the other member being complexed with alkaline phosphatase or other suitable enzyme as a trigger. The principal restriction on the identity of the binding pairs is that the first binding pair be distinct from, and not interact with, the second binding pair.

In addition to alkaline phosphatase, other enzymes which may be used to cleave the enzymatically cleavable group from the dioxetane include: acid phosphatases, esterases, decarboxylases, phospholipase D, β -xylosidase, β -D-fructosidase, thioglucosidase, β -D-galactosidase, α -D-galactosidase, α -D-glucosidase, β -D-glucosidase, α -D-mannosidase, β -D-mannosidase, β -D-fructofuranoside, β -D-glucosiduronase, and trypsin.

In addition to being of particular interest as organic moieties that constitute diagnostic markers, protease enzymes are also of considerable interest as enzyme labels. Examples of diagnostic protease markers include cathepsin B (cancer), cathepsin G (emphysema, rheumatoid arthritis, inflammation), plasminogen activator (thrombosis, chronic inflammation, cancer) and urokinase (cancer). Assays for protease detection are therefore needed to monitor protein stability in various biological and commercial processes.

In an alternative embodiment of this assay a homogeneous chemiluminescent energy transfer assay is provided. In this approach, one end of the cleavage sequence peptide bears

a 1,2-dioxetane or dioxetane precursor. Direct attachment of this moiety, followed by oxygenation to form the dioxetane, is enabled in U.S. Patent Application Serial No. 08/767,479, allowed and incorporated herein by reference. The other end of this peptide bears an energy accepting fluorescent molecule such as fluorescein, or any of a variety of similar fluorescing moieties, such as those disclosed in U.S. Patent 5,004,565 and 5,208,148 which are incorporated herein by reference. The peptide is sufficiently short (no more than about 10 amino acid residues) such that the dioxetane is in close physical association with the fluorescent label. Upon triggering of the dioxetane, which can be effected by addition of an enzyme, or pH alteration, or application of heat or other triggers, the dioxetane decomposes, emitting energy which excites the fluorescent moiety which then fluoresces if no cleavage has occurred (a positive test for protease inhibition). If cleavage has occurred, the dioxetane and fluorescent moieties are no longer in close physical relationship, and the light is emitted by the chemiluminescent dioxetane. The wavelength of the fluorescent emitter is characteristically shifted markedly from that of the dioxetane, allowing easy discrimination in a homogenous assay.

Another approach to a homogeneous assay using an enol ether or dioxetane label is to use a quenching group in place of the fluorescent acceptor. In this format, upon triggering of the dioxetane label, there will be no light from the intact peptide due to the quenching group. If cleavage of the peptide substrate by a protease occurs prior to triggering, emission from the dioxetane group will be observed upon triggering. This quenching process is subject to the same distance restraints as the energy transfer process described above. A suitable quenching group for the substrate shown in Figure 14 would be a dabcyI group in place of the fluorescein group. Any non-fluorescent light absorbing

dye, which has an absorption spectra which overlaps the emission spectra of the donor dioxetane could be used as a quenching group.

Variations on this approach can be made for many types of proteases. Direct covalent or hydrophobic attachment of a peptide substrate to a microwell surface may be required for some proteases. Alternatively, direct synthesis of the peptide on the solid phase may also have advantages.

The heterogenous assay of this invention has been exemplified, below, by reference to an assay adapted to detect inhibition of HIV-1 protease. This example is demonstrative only, and not intended to be limiting.

Materials and Methods

Reagents:

1. The HIV FLAG peptide was custom synthesized by Genemed Synthesis, Inc.
2. Anti-FLAG MI Monoclonal Antibody, FLAG Octapeptide, and Enterokinase were purchased from Eastman Kodak Company.

Neutravidin coated plates: Neutravidin™ Biotin Binding Protein (PIERCE) was diluted to 5ug/ml or 0.1 ug/ml in BupH™ Carbonate-Bicarbonate buffer (PIERCE), 1 mg/ml BSA (fraction V Sigma) and 100ul added to white 96 well plates (Dynatech-Microlite1) and incubated for 2 hours at 37 °C. The wells were then washed with PBS, 0.1%(v/v) Tween 20 and blocked with PBS, 0.1%(V/V) Tween 20, 1 mg/ml BSA overnight at 4 degrees.

Protease assays: Rec HIV-PR1 (affinity purified -BACHEM) was diluted in 0.1M NaAcetate, 1M NaCl, 1mM EDTA, 1mM DTT, 1 mg/ml BSA, pH 4.7 or pH 5.5 and added to Neutravidin coated plates. Rec HIV-1 inhibitor (Acetyl-pepstatin(BACHEM)), compounds from plate-1001 (Sigma Drug Standards) and/or DMSO was added followed

with a 5 minute preincubation before the addition of HIV-1 peptides I or II, (FAM-spacer-Ser-Gln-Asn-Tyr-Pro-Ile-Val-Gln-spacer-Biotin (Perkin-Elmer) or (FLAG-Ser-Nle-Ala-Glu-Phe-Leu-Val-Arg-Ala-Lys-His-Spacer-Biotin) or DABCYL-y-Abu-Ser-Gln-Asn-Tyr-Pro-Ile-Val-Gln-EDANS (BACHEM) (prior art) respectively. Reactions were done at ambient and/or 37°C. For HPLC analysis, reactions were stopped with 0.1 % TFA and placed on ice before injection. Assays with the BACHEM fluorogenic substrate were measured directly on a LS50B fluorescence spectrophotometer (Perkin-Elmer). Neutravidin coated assay plates with captured biotinylated-fluorescein labeled peptide I were washed with PBS, 0.1 % (V/V) Tween20, 1 mg/ml BSA and incubated for 60 min. with anti-fluorescein-alkaline phosphatase Fab fragments (Boehringer Mannheim) at ambient temperature. Plates were washed again with PBS buffer followed by a 10mM Tris-HCL, 1mM MgCl₂ pH 9.8 wash. 100ul of CSPD/Sapphire II™ ready to use alkaline phosphatase was added and emitted light was measured in a TR717® microplate luminometer (Tropix).

HPLC analysis: Analytical HPLC on the HIV-1 protease substrate cleavage products was done on a Perkin-Elmer UV/Vis detector with series 200lc pump, Perkin-Elmer pecosphere 5 C18, 5um, 4.6mm x 15cm, gradient of water(0.1 % TFA) and CH₃CN (0.1 % TFA) from 30% to 60% over 10 minutes at 1 ml/min.

Additions and dilutions were made with a Zymark RapidPlate-96 pipeting station (Zymark Corp.) and washes were done with a Tecan 96PW washer (Tecan).

The present inventors have successfully developed an automated screening assay as shown below to search for novel HIV-1 protease inhibitors.

AUTOMATION STEPS

#	DESCRIPTION	VOLUME (μ L)	AUTOMATION COMPONENT	TIME (MIN)
1	Add protease buffer containing 50nM HIV-1 protease	50 μ L	Zymark Rapid plate 96-well pipettor	1 min
2	Add test compound	5 μ L	Zymark Rapid plate 96-well pipettor	1 min
3	Add HIV Perkin Elmer peptide @ 444nM	45 μ L	Zymark Rapid plate 96-well pipettor	1 min
4	Incubate (32°C)			60 min
5	Wash	3X100 μ L	Tecan 96PW washer	0.75 min
6	Add Anti-Fluorescein-AP conjugate	100 μ L	Zymark Rapid plate 96-well pipettor	1 min
7	Incubate (Room Temp.)			30-60 min
8	Wash	5X100 μ L	Tecan 96PW washer	1 min
9	Add CSPD + Sapphire II	100 μ L	Zymark Rapid plate 96-well pipettor	1 min
10	Read plate		Tropix TR717 Luminometer	30 min + 1.4 min

We have shown that this protease assay can be set up as an endpoint capture assay with minimal reagent use. An HIV-1 peptide substrate I concentration of 200nM and a rec HIV-1 protease concentration of 25-50nM have been shown to be optimal working concentrations with good sensitivity and acceptable signal:noise. A resonance energy transfer assay reported by Abbott Laboratories, Matayoshi, E.D. et al., Science 247: 954-958 (1990), utilizes micromolar amounts of HIV-1 peptide II. This assay is reported as an assay with a linear detection window of less than 5 minutes, which we have found to have a poor signal to noise ratio making it a difficult robotic assay to automate. Figure 6. This assay gave a maximum s/n ratio of 4, while the inventive assay gives a s/n ratio of greater

than 140 when using the FLAG/anti-FLAG ligand binding pair. By utilizing common laboratory reagents such as biotin and fluorescein one can eliminate the need to generate sequence specific monoclonal antibodies as described in the alternative HIV peptide capture based assay by Fournout, S. et al., Anal. Chem. 69, 1746-1752 (1997). We have shown that DMSO is well tolerated in the assay. The IC_{50} of acetyl-pepstatin in our capture assay was 2-3uM compared to 0.3uM reported by Fournout, S. et al.

This invention has been disclosed in terms of both generic description and specific example. Variations will occur to those of ordinary skill in the art, including peptide moiety identities, specific proteolytic enzymes to be employed, enhancement agents and enhancement additives, and specific assay formats without the exercise of inventive faculty. Such variations remain within the scope of the invention, save for variations excluded by the recitation of the claims presented below.

CLAIMS:

1. A method for conducting an assay to determine whether a target compound exhibits activity as a protease inhibitor, comprising:

combining said protease and said target compound in an environment further comprising a construct, wherein said environment is such that said protease cleaves said construct in the absence of inhibition of said protease,

said construct comprising an amino acid sequence terminating, at a first end, with a first member of a first ligand binding pair and at a second end with a first member of a second ligand binding pair, wherein

said first member of said first ligand binding pair binds to a second member of said first ligand binding pair which second member is bound to a surface,

removing any fragments of said construct cleaved by said protease,

adding a complex of a second member of said second ligand binding pair complexed with an enzyme and allowing said complex to bind to any said first member of said second ligand binding pair present,

removing any unbound complex,

adding a 1,2-dioxetane which is a substrate for said enzyme and observing any chemiluminescence released thereby, wherein emission of said chemiluminescence is indicative of protease inhibition activity by said target compound.

2. The method of Claim 1, wherein said chemiluminescence observed is measured, and wherein the amount of chemiluminescence observed is correlated with the degree of inhibitory activity exhibited by said compound.

3. The method of Claim 1, wherein said assay is a transfer free single plate endpoint assay.
4. The method of Claim 1, wherein said surface is at least a portion of at least one well of a test plate.
5. The method of Claim 1, wherein said second member of said second ligand binding pair is an antibody.
6. The method of Claim 1, wherein said first ligand binding pair is biotin and a compound which binds to biotin at least as tightly as avidin.
7. The method of Claim 1, wherein said second ligand binding pair is biotin and a compound which binds to biotin at least as tightly as avidin.
8. The method of Claim 6, wherein said compound which binds to biotin at least as tightly as avidin is avidin or strepavidin.
9. The method of Claim 7, wherein said compound which binds as tightly as avidin is avidin or strepavidin.
10. The method of Claim 1, wherein said first member of said first ligand binding pair is biotin and said first member of said second ligand binding pair is fluorescein.
11. The method of Claim 1, wherein said second ligand binding pair is FLAG and an antibody therefor.
12. The method of Claim 1, wherein said protease is selected from the group consisting of a serine protease, a cysteine protease, an aspartic protease and a metalloproteinase.
13. The method of Claim 1, wherein said protease is selected from the group consisting of HIV-1 protease, caspases, cathhepsins, hydrolase, L-proteinase, calpain,

interleukin converting proteases, urokinase, trypsin, thrombin, HIV-2 protease, Yapsin I, Yapsin 3, Plasmepsin I, Plasmepsin II, collagenase, gelatinases, stromelysin, amino peptidase and elastase.

14. The method of Claim 13, wherein said protease is HIV-1 protease.

15. A protease substrate for conducting an assay to determine protease inhibitory activity in a target compound, said substrate comprising:

a polypeptide, said polypeptide including a cleavage site specific for said protease, said polypeptide bearing at a first end a first member of a first ligand binding pair and at another end a first member of a second ligand binding pair, wherein said first member of said first ligand binding pair and said first member of said second ligand binding pair do not bind to each other.

16. The substrate of Claim 15, wherein said first and second ligand binding pair is comprised, independently, of an antigen and an antibody therefore, or biotin and a compound which binds as tightly to biotin as avidin.

17. The substrate of Claim 15, wherein said substrate has the sequence Fluorescein-Spacer-Ser-Glu-Asu-Tyr-Pro-Ile-Val-Glu-Spacer-Biotin, or FLAG-Ser-Nle-Ala-Glu-Phe-Leu-Val-Arg-Ala-Hys-His-Spacer-Biotin.

18. A homogenous assay for detection of protease inhibitory activity in a target compound, comprising:

combining said protease and said target compound in an environment further comprising a construct, wherein said environment is such that said protease cleaves said construct in the absence of inhibition of said protease, said construct comprising a polypeptide of 2-10 amino acids, said polypeptide terminating, at a first end, and a moiety

which is a 1,2-dioxetane or precursor thereof which can be oxygenated in said environment to provide a 1,2-dioxetane moiety and, at a second end, with an energy accepting fluorescent moiety,

oxygenating said precursor if present to form a 1,2-dioxetane moiety,

causing said 1,2-dioxetane moiety to decompose, and observing the wavelength of light emitted from said environment, wherein said wavelength, if the wavelength of said fluorescent emitter, is indicative of inhibitory activity, and said wavelength, if the wavelength of said dioxetane, is indicative of absence of inhibitory activity.

19. A construct for conducting a homogenous assay to determine protease inhibitory activity in a target compound, comprising:

a polypeptide of 2-10 amino acid residues, wherein said polypeptide includes a cleavage site specific for said protease, said polypeptide bearing, at one end, a fluorescent energy accepting moiety or a chemiluminescence quenching moiety and at another end, a 1,2-dioxetane or 1,2-dioxetane precursor moiety that can be oxygenated to form a 1,2-dioxetane moiety in the course of said assay.


20. A kit for conducting an assay to determine whether a protease inhibitor is present in a sample, comprising in one or more containers:

- (a) a peptide which is a substrate for said protease, wherein said peptide substrate is labeled with (i) a first member of a first ligand binding pair, and (ii) a first member of a second ligand binding pair;
- (b) a chemiluminescent 1,2-dioxetane substrate containing an enzymatically cleavable group, which substrate is capable of producing light in the presence of

an enzyme which cleaves said enzymatically cleavable group from said substrate.


21. The kit of Claim 20, wherein said protease is HIV-1.
22. The kit of Claim 21, wherein wherein said peptide substrate comprises the amino acid sequence: -Ser-Gln-Asn-Try-Pro-Ile-Val-Gln-.
23. The kit of Claim 20, wherein said first member of said second ligand binding pair is fluorescein.
24. The kit of Claim 20, wherein said first member of said first ligand binding pair is biotin.
25. The kit of Claim 20, additionally comprising a polymeric quaternary onium salt enhancement agent for said 1,2-dioxetane.
26. The kit of Claim 25, wherein said enhancement agent comprises polyvinylbenzyltributyl ammonium chloride.
27. The kit of Claim 20, further comprising a enzyme complexed with a second member of said second ligand binding pair.
28. The kit of Claim 27, wherein said enzyme is alkaline phosphatase.

1/16




2/16

NO INHIBITION

**FIG. 2A**

INHIBITION WITH ACETYL-PEPSTATIN

**FIG. 2B**

SUBSTITUTE SHEET (RULE 26)

3/16




FIG. 3

4/16




FIG. 4

5/16












FIG. 5A

6/16



7/16







FIG. 6

8/16

*FIG. 7A**FIG. 7B*

9/16

*FIG. 8*

10/16






FIG. 9

11/16

*FIG. 10A**FIG. 10B*

SUBSTITUTE SHEET (RULE 26)

12/16






FIG. 11




REC.HIV-1 (nM) AND CONDITION

FIG. 12

14/16

*FIG. 13*

*FIG. 14A*




FIG. 14B

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/03929**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :C12Q 1/37

US CL :435/23, 975

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/23, 975

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Computer Search - CAPLUS, MEDLINE, APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,605,795 A (BRONSTEIN) 25 FEBRUARY 1997, see entire document.	1-28
A	US 5,591,591 A (BRONSTEIN et al.) 07 JANUARY 1997, see entire document.	1-28
A	AYYAGARI, M.S., et al. Chemiluminescence-Based Inhibition Kinetics Of Alkaline Phosphatase In The Development Of A Pesticide Biosensor, Biotechnol. Prog. 1995, Vol. 11, pages 699-703, see entire document.	1-28
A	MATAYOSHI, E.D., et al. Novel Fluorogenic Substrates For Assaying Retroviral Proteases By Resonance Energy Transfer. Science. 23 FEBRUARY 1990, Vol.247, pages 954-958, see entire document.	1-28



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

19 MAY 1998

Date of mailing of the international search report

12 JUN 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CHARLES PATTERSON

Telephone No. (703) 308-0196